

Adverse events should be reported. Reporting forms and information can be found at <https://yellowcard.mhra.gov.uk/>

**Adverse events should also be reported to
McNeil Products Limited on freephone 0808 238 9999.**

DAKTARIN Sugar Free 2% Oral Gel (MICONAZOLE 20mg per gram) Product Information

Presentation:

White gel with an orange taste containing 20mg miconazole per gram.

Indication:

Oral treatment of superficial fungal (e.g. Candida) infections of the oropharynx in adults and paediatric patients 4 months and older.

Dosage and Administration:

For topical treatment of the oropharynx. Infants: 4 – 24 months: 1.25 ml ($\frac{1}{4}$ measuring spoon of gel), applied four times a day after meals. Each dose should be divided into smaller portions and the gel should be applied to the affected area(s) with a clean finger. The gel should not be applied to the back of the throat due to possible choking. The gel should not be swallowed immediately but kept in the mouth as long as possible. Adults and children 2 years of age and older: 2.5ml ($\frac{1}{2}$ measuring spoon of gel), applied four times a day after meals. The gel should not be swallowed immediately but kept in the mouth as long as possible. The treatment should be continued for at least a week after the symptoms have disappeared. For oral candidosis, dental prostheses should be removed at night and brushed with the gel.

Contraindications:

Hypersensitivity to the active substance(s), other imidazole derivatives or to any of the excipients listed in SPC section 6.1. Infants less than 4 months of age or in those whose swallowing reflex is not yet sufficiently developed. Liver dysfunction. DAKTARIN Sugar Free 2% Oral Gel should not be given concurrently with astemizole, cisapride, dofetilide, mizolastine, pimozide, quinidine, sertindole, terfenadine, ergot alkaloids, triazolam, oral midazolam, warfarin or HMG-CoA reductase inhibitors such as simvastatin and lovastatin.

Precautions:

If the concomitant use of Daktarin and drugs such as anticoagulants like warfarin, is planned, the anticoagulant effect should be carefully monitored and titrated. Miconazole is systemically absorbed and is known to inhibit CYP2C9 and CYP3A4 which can lead to prolonged effects of warfarin. Bleeding events, some with fatal outcomes have been reported with concurrent use of miconazole oral gel and warfarin. It is advisable to monitor miconazole and phenytoin levels, if these two drugs are used concomitantly. In patients using certain oral hypoglycaemics such as sulphonylureas, an enhanced therapeutic effect leading to hypoglycaemia may occur during concomitant treatment with miconazole and appropriate measures should be considered. Caution should be used when Daktarin is taken concomitantly with drugs subject to metabolism by CYP3A4 and CYP2C9. The lower age limit should be increased to 5 – 6 months of age

for infants who are pre-term, or infants exhibiting slow neuromuscular development. Choking in infants and young children - Particularly aged 4 months – 2 years, caution is required, to ensure that the gel does not obstruct the throat. Hence, the gel should not be applied to the back of the throat. Observe the patient for possible choking. This medicine contains 7.85 mg of alcohol (ethanol) in each gram of oral gel. The amount in single maximum dose of this medicine for an adult (10 ml oral gel) is equivalent to less than 3 ml beer or 1 ml wine. The small amount of alcohol in this medicine will not have any noticeable effects. This medicinal product contains orange flavour (containing: citral, citronellol, linalool, geraniol, d-limonene) and cocoa flavour (containing: benzyl alcohol, benzyl benzoate) that may cause allergic reactions. Benzyl benzoate may cause mild local irritation. Benzyl alcohol may cause allergic reactions or may cause mild local irritation. Caution should be exercised in administration of benzyl alcohol due to the following side effects in these populations: gasping syndrome in children and metabolic acidosis for breastfeeding and pregnant women or patients with kidney disease.

Fertility, Pregnancy and Lactation:

Daktarin Sugar Free 2% Oral Gel should be avoided in pregnant women if possible. Consult a doctor prior to use. The potential hazards should be balanced against the possible benefits. Caution in nursing mothers.

Side effects:

Increases in INR and bleeding events such as epistaxis, contusion, haematuria, melaena, haematemesis, haematoma and haemorrhages have been reported in patients treated with oral anticoagulants such as warfarin in association with miconazole oral gel. Some events had fatal outcomes.

Common: dry mouth, nausea, oral discomfort, vomiting, regurgitation and abnormal product taste.

Uncommon: dysgeusia.

Unknown frequency: anaphylactic reaction, hypersensitivity, choking, diarrhoea, stomatitis, tongue discolouration, hepatitis, angioedema, toxic epidermal necrolysis, Stevens-Johnson syndrome, urticaria and rash. Acute generalized exanthematous pustulosis, drug reaction with eosinophilia and systemic symptoms. See SPC for details.

RRP (ex-VAT): 15g tube £7.48

Legal category: P.

PL Holder: McNeil Products Ltd, 50 -100 Holmers Farm Way, High Wycombe, Buckinghamshire, HP12 4EG.

PL Number(s): 15513/0296.

Date of prep: 01 March 2023.